

MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A148 432

12

OFFICE OF NAVAL RESEARCH

Contract No. N00014-83-K-0611

Task No. NR 053-842

TECHNICAL REPORT NO. UK/DC/TR-3 ✓

Syntheses and Reactions of Pyrazaboles

by

W. J. Layton, K. Niedenzu,\* P. M. Niedenzu and S. Trofimenko

Prepared for publication in

INORGANIC CHEMISTRY

University of Kentucky  
Department of Chemistry  
Lexington, KY 40506

November 1984

DTIC  
ELECTE  
DEC 12 1984  
S E D

DTIC FILE COPY

Reproduction in whole or in part is  
permitted for any purpose of the United States Government

This document has been approved for public  
release and sale; its distribution is unlimited

11/2/84

unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER UK/DC/TR-3	2. GOVT ACCESSION NO. AD-A148432	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle)  SYNTHESES AND REACTIONS OF PYRAZABOLES		5. TYPE OF REPORT & PERIOD COVERED Interim Technical Report, 1984
		6. PERFORMING ORG. REPORT NUMBER 3
7. AUTHOR(s) W. J. Layton, K. Niedenzu, P. M. Niedenzu and S. Trofimenko* (*E. I. DuPont de Nemours & Co.)		8. CONTRACT OR GRANT NUMBER(s)  N00014-83-K-0611
9. PERFORMING ORGANIZATION NAME AND ADDRESS Univeristy of Kentucky, Department of Chemistry Lexington, KY 40506-0055		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS  NR 053-842
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arlington, VA 22217		12. REPORT DATE November 1984
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 22
		15. SECURITY CLASS. (of this report)  Unclassified
15a. DECLASSIFICATION/DOWNGRADING SCHEDULE		
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)  Prepared for publication in INORGANIC CHEMISTRY		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  pyrazaboles NMR amine-boranes		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  The unsymmetrical pyrazabole $(C_6H_5)_2B(pz)_2BH_2$ ( $pz = N_2CH_3 =$ pyrazolyl) was prepared by the reaction of $K[(C_6H_5)_2Bpz_2]$ with $(CH_3)_2N_2BH_2$ ; subsequent halogenation with $Br_2$ yielded $(C_6H_5)_2B(pz)_2BBR_2$ . Similarly, $(C_2H_5)_2B(pz')_2BH_2$ ( $Hpz' =$ 3,5-dimethylpyrazole) was converted to		

DD FORM 1473  
1 JAN 73EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-LF-014-6601

unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

$(C_2H_5)_2B(\mu-pz')_2EBr_2$ . Reaction of  $H_2B(\mu-pz')_2BH_2$  with (even an excess of)  $BBr_3$  gave a mixture of cis and trans isomers of  $HBrB(\mu-pz')_2BBr$ , whereas reaction with  $Br_2$  afforded  $Br_2B(\mu-pz')_2BBr_2$ . Reaction of the latter compound with Kpz yielded  $pz_2B(\mu-pz')_2Bpz_2$ , the first characterized pyrazolylpyrazabole containing different pyrazolyl moieties bonded to the same boron atom. A second, polymeric modification of  $Hpz'B(\mu-pz')_2BHpz'$  was identified; it appears to be the one reacting with additional  $Hpz'$  to form  $pz'_2B(\mu-pz')_2Bpz'_2$ . The latter forms a monohydrate in a reversible reaction, but no similar interaction occurs with  $NH_3$ .

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Avail and/or	
Dist	Special
A-1	



unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Contribution from the Department of Chemistry, University of Kentucky, Lexington, Kentucky 40506; and the Polymer Products Department, E. I. DuPont de Nemours & Co., Parkersburg, West Virginia 26101

Syntheses and Reactions of Pyrazaboles<sup>1</sup>

W. J. Layton,<sup>a</sup> K. Niedenzu,<sup>a</sup> P. M. Niedenzu,<sup>a</sup> and S. Trofimenko<sup>b</sup>


-----  
<sup>a</sup> University of Kentucky; <sup>b</sup> E. I. DuPont de Nemours & Co.  
-----

Received.....

The unsymmetrical pyrazabole  $(C_6H_5)_2B(\mu\text{-pz})_2BH_2$  ( $\text{pz} = N_2C_3H_3 =$  pyrazolyl) was prepared by the reaction of  $K[(C_6H_5)_2Bpz_2]$  with  $(CH_3)_3NBH_2I$ ; subsequent halogenation with  $Br_2$  yielded  $(C_6H_5)_2B(\mu\text{-pz})_2BBr_2$ . Similarly,  $(C_2H_5)_2B(\mu\text{-pz}')_2BH_2$  ( $\text{Hpz}' =$  3,5-dimethylpyrazole) was converted to  $(C_2H_5)_2B(\mu\text{-pz}')_2BBr_2$ . Reaction of  $H_2B(\mu\text{-pz}')_2BH_2$  with (even an excess of)  $BBr_3$  gave a mixture of cis and trans isomers of  $HBrB(\mu\text{-pz}')_2BBr$ , whereas reaction with  $Br_2$  afforded  $Br_2B(\mu\text{-pz}')_2BBr_2$ . Reaction of the latter compound with  $Kpz$  yielded  $pz_2B(\mu\text{-pz}')_2Bpz_2$ , the first characterized pyrazolylpyrazabole containing different pyrazolyl moieties bonded to the same boron atom. A second, polymeric modification of  $\text{Hpz}'B(\mu\text{-pz}')_2B\text{Hpz}'$  was identified; it appears to be the one reacting with additional  $\text{Hpz}'$  to form  $pz'_2B(\mu\text{-pz}')_2Bpz'_2$ . The latter forms a monohydrate in a reversible reaction, but no similar interaction occurs with  $NH_3$ .

### Introduction

The pyrazaboles of the general structure 1 (with positional numbering) are a chemically and thermally remarkably stable class of heterocycles containing four-coordinate boron.<sup>2</sup>



1

Surprisingly, their chemistry has been studied only to a relatively small extent. However, the discovery of isolable symmetrical cleavage products of pyrazaboles, i.e., pyrazol-1-ylboranes containing trigonal boron,<sup>3</sup> has led to a renewed interest in pyrazabole chemistry and brought about new features, especially with respect to electrophilic attack at B-H bonds of pyrazaboles.<sup>4</sup> Moreover, it was found that methylation at the 1,3,5 and 7 sites of a pyrazabole skeleton imposes significant differences in chemical behavior.<sup>5</sup>

The present work is concerned with a further exploration of syntheses and the reactivity of pyrazaboles.

### Experimental Section

Elemental analyses were performed by the Schwarzkopf Microanalytical Laboratory, Woodside, NY. Melting points (uncorr.) were determined on a Mel-Temp block.

Mass spectral data (unless otherwise noted: 70 eV) were obtained from the University of Kentucky Mass Spectrometry Center and were recorded on a PE-Hitachi RMU-7 or VG ZAB-2F instrument. Data are listed for ions of 5% or higher relative abundances only. NMR spectra were recorded on a Varian XL-200 spectrometer. Chemical shift data are given in ppm with positive values indicating downfield from the reference (internal  $\text{Me}_4\text{Si}$  for  $^1\text{H}$  and  $^{13}\text{C}$ , external  $\text{Et}_2\text{OBF}_3$  for  $^{11}\text{B}$ ). Abbreviations: s = singlet, d = doublet, t = triplet, q = quartet, p = quintet, m = unresolved multiplet. An asterisk denotes a broad signal. Coupling constants J are given in Hz. Details for HOMCOR and HETCOR NMR experiments have been given elsewhere.<sup>6</sup> Infrared spectra were recorded on a PE Model 621 instrument under standard operating conditions.

Ammine-triphenylborane. To 2.5 L of commercially available (E. I. DuPont de Nemours & Co.) aqueous solution of  $\text{Na}[\text{HOB}(\text{C}_6\text{H}_5)_3]$  (ca. 7% plus 3% excess NaOH) was added 1 L of conc. aqueous ammonia solution (large excess). The resultant colorless precipitate was collected, washed with water until the washings were no longer basic, and pressed dry. The still moist material was stirred with 1.2 L ethyl acetate. The lower (aqueous) layer of the resultant mixture was discarded. The upper (organic)



layer was stirred with Celite and anhydrous  $\text{MgSO}_4$  and filtered. The filtrate was concentrated until a thick paste remained. Sufficient toluene was added to make a stirrable slurry. The desired compound was collected as fine needles that were dried in vacuum (ca. 80 % yield); it begins to decompose near  $180^\circ\text{C}$  and is completely molten/decomposed near  $225^\circ\text{C}$ . Anal. Calcd. for  $\text{C}_{18}\text{H}_{18}\text{BN}$  (259.16): C, 83.42; H, 7.00; B, 4.17; N, 5.40. Found: C, 83.20; H, 7.29; B, 3.98; N, 5.29.  $\delta(^{11}\text{B})$  (solution in  $\text{CDCl}_3$ ) =  $-2.7$  ( $h_{\frac{1}{2}} = 200$  Hz).

Potassium diphenyl-dipyrazol-1-yl-borate. A 500-mL flask equipped with a small Vigreux column was charged with 150 g (2.2 mole) of pyrazole and 22.4 g (0.4 mole) KOH pellets. The mixture was heated until all water distilled out and pyrazole started to distil. Heating was stopped immediately and the liquid was allowed to cool to  $110\text{--}120^\circ\text{C}$ . Then 103.5 g (0.4 mole) of  $\text{H}_3\text{NB}(\text{C}_6\text{H}_5)_3$  was added in small portions, and heating was resumed at such a rate that benzene distilled out slowly. A total of 35 mL was collected. The melt was then poured into 600 mL of rapidly stirred toluene. The precipitated  $\text{K}[(\text{C}_6\text{H}_5)_2\text{Bpz}_2]$  was collected and washed with hot toluene and was then dried under vacuum to yield 101 g (75 %) of the desired compound. An additional 15 g (11 %) of slightly less pure material could be recovered from the filtrate (excess pyrazole being removed by distillation). Anal. Calcd. for  $\text{C}_{18}\text{H}_{16}\text{BKN}_4$  (338.26): C, 63.91; H, 4.77; B, 3.20; K, 11.56; N, 16.56. Found: C, 63.88; H, 5.10; B, 2.94; N, 16.89.

NMR data (solution in DMSO- $d_6$ ):  $\delta(^1\text{H}) = 7.43$  (1H, d,  $J = 2.4$ ), 7.16 (1H, unresolved), 7.03 + 7.00 (4H, unresolved), 5.99 (1H, unsymm. t = 2 overlapping d,  $J$  ca. 2);  $\delta(^{11}\text{B}) = 1.3$  ( $h_{\frac{1}{2}} = 325$  Hz);  $\delta(^{13}\text{C}) = 153.9^*$ , 137.8 (d,  $J = 178$ , of t,  $J = 7$ ), 134.0 (d,  $J = 157$ ), 133.7 (d,  $J = 192$ ), 125.6 (d,  $J = 155$ , of d,  $J = 5$ ), 124.0 (d,  $J = 157$ , of t,  $J = 7$ ), 101.5 (d,  $J = 171$ , of t,  $J = 11$ ). Solution in  $\text{D}_2\text{O}$ :  $\delta(^{11}\text{B}) = 0.5$  ( $h_{\frac{1}{2}} = 260$  Hz);  $\delta(^{13}\text{C})$  (proton-decoupled) = 133.1, 130.6, 126.2, 124.2, 104.1.

Trimethylamine-monoiodoborane.<sup>8</sup> NMR data (solution in  $\text{CDCl}_3$ ):  $\delta(^1\text{H}) = 2.87$  (9H, s), ca. 2.6 (2H, very very broad);  $\delta(^{11}\text{B}) = -9.5$  (t,  $J = 131$ ). Literature data:  $\delta(^1\text{H})$  (in  $\text{CH}_2\text{Cl}_2$ ) = 2.82;<sup>9</sup>  $\delta(^{11}\text{B})$  (in benzene) = -8.9 (t,  $J = 142$ ).<sup>10</sup>

4,4-Diphenylpyrazabole. A mixture of 18.7 g (55 mmole) potassium diphenyldipyrazol-1-ylborate, 11.3 g (57 mmole) of trimethylamine-monoiodoborane<sup>8</sup> and 300 mL toluene was stirred and slowly heated to 80°C. This temperature was maintained for 6 h and was then raised to reflux for 35 h. The hot mixture was filtered and the solvent was evaporated. The remaining crystals were crushed and washed with two 50-mL portions of water. After drying, they were stirred twice with 50-mL portions of petroleum ether. The crystals were then dissolved in chloroform and the clear filtrate was evaporated to give 14.62 g (82 %) of the crude product, mp 158-162°C. After recrystallization from toluene the compound melted at 168-170°C. Anal. Calcd. for  $\text{C}_{18}\text{H}_{18}\text{B}_2\text{N}_4$  (311.99): C, 69.30; H, 5.82; B, 6.93; N, 17.96. Found: C, 69.06; H, 5.93; B, 7.08; N, 17.82.

NMR data (solution in  $\text{CDCl}_3$ ):  $\delta(^1\text{H}) = 7.66$  (1H, d,  $J = 2.0$ ), 7.37 (1H, d,  $J = 2.3$ ), 7.24-7.21 (3H, m), 6.99-6.94 (2H, m), 6.31 (1H, t,  $J = 2.2$ ), 3.4\* (2H);  $\delta(^{11}\text{B})$  (proton-decoupled) = +1.7 (1B,  $h_{\frac{1}{2}} = 610$  Hz), -8.4 (1B,  $h_{\frac{1}{2}} = 220$  Hz; broadens in the proton-coupled spectrum but could not be resolved);  $\delta(^{13}\text{C}) = 145^*$ , 136.3 (d,  $J = 190$ , of t,  $J = 7$ ), 135.7 (d,  $J = 189$ , of t,  $J = 7$ ), 133.6 (d,  $J = 156$ , of unresolved m), 127.3 (d,  $J = 161$ , of unresolved m), 126.8 (d,  $J = 159$ , of t,  $J = 8$ ), 105.3 (d,  $J = 180$ , of t,  $J = 9$ ).

Pyrazabole.<sup>7</sup>  $\delta(^{11}\text{B})$  (solution in  $\text{DMSO-d}_6$ ) = -8.2 ( $h_{\frac{1}{2}} = 250$  Hz, proton-decoupled;  $h_{\frac{1}{2}} = 350$  Hz, proton-coupled). Additional NMR data have been reported elsewhere.<sup>6</sup> - First ionization potential: 12.24 eV (from CNDO calculations estimated by Koopmans theorem);  $9.0 \pm 0.2$  eV (from mass spectrometry).

4,4,8,8-Tetraphenylpyrazabole.<sup>4</sup> NMR data (solution in  $\text{CDCl}_3$ ):  $\delta(^1\text{H}) = 7.48$  (2H, d,  $J = 2.5$ ), 7.03 + 7.00 (3H, m), 6.81 + 6.78 (2H, m), 6.41 (1H, t,  $J = 2.5$ );  $\delta(^{11}\text{B}) = +1.7$  ( $h_{\frac{1}{2}} = 320$  Hz);  $\delta(^{13}\text{C}) = 145.5^*$ , 137.3 (d,  $J = 190$ , of t,  $J = 7$ ), 133.2 (d,  $J = 154$ , of m), 127.0 (d,  $J = 157$ , of m), 126.4 (d,  $J = 159$ , of t,  $J = 7.5$ ), 105.5 (d,  $J = 181$ , of t,  $J = 8.5$ ). Suggested assignments (based on HOMO-COR and HETCOR experiments and on coupling constant data):  $\delta(^1\text{H})/\delta(^{13}\text{C}) = 7.03 + 7.00/126.4$ , 127.0, o- and p-phenyl; = 6.80/133.2, m-phenyl; = 7.48/137.3, N-bonded CH of pyrazolyl groups; = 6.41/105.5, central CH of pyrazolyl groups.

4,4-Diphenyl-8,8-dibromopyrazabole. A solution of  $\text{Br}_2$  in  $\text{CH}_2\text{Br}_2$  was slowly added to a solution of 3.12 g (10 mmol) of 4,4-diphenylpyrazabole in  $\text{CH}_2\text{Br}_2$  until the color of free  $\text{Br}_2$  remained. The mixture was refluxed for 15 min and solvent was stripped off under reduced pressure until a volume of ca. 30 mL remained. On cooling to room temperature, 3.0 g (64%) of the desired compound precipitated. The latter was collected, washed with 5 mL portions each of toluene, chloroform and petroleum ether. The resultant colorless crystals had a mp  $220^\circ\text{C}$  (after drying in vacuum at  $60^\circ\text{C}$ ). Anal. Calcd. for  $\text{C}_{18}\text{H}_{16}\text{Br}_2\text{N}_4$  (469.78): C, 46.02; H, 3.43; Br, 34.02; N, 11.93. Found: C, 45.53; H, 3.68; Br, 33.61; N, 11.76.

NMR data (solution in  $\text{CDCl}_3$ ):  $\delta(^1\text{H}) = 8.48$  (1H, d,  $J = 2.5$ ), 7.57 (1H, d,  $J = 2.4$ ), 7.28-7.21 (3H, m), 6.98-6.94 (2H, m), 6.61 (unsymmetrical t,  $J = 2.5$ );  $\delta(^{11}\text{B}) = +1.9$  ( $h_{\frac{1}{2}} = 300$  Hz), -6.6 ( $h_{\frac{1}{2}} = 40$  Hz), area ratio 1:1;  $\delta(^{13}\text{C})$  (proton-decoupled) = 145\*, 139.5, 139.1, 133.2, 127.6, 127.5, 107.9.

1,3,5,7-Tetramethyl-4,4-diethyl-8,8-dibromopyrazabole. A solution of  $\text{Br}_2$  in  $\text{CH}_2\text{Br}_2$  was added dropwise with stirring to a solution of 3.37 g (12.5 mmole) of 1,3,5,7-tetramethyl-4,4-diethylpyrazabole<sup>11</sup> in  $\text{CH}_2\text{Br}_2$ . Once the color of free bromine remained, the mixture was refluxed for 15 min. The volume of the solution was then reduced under vacuum to ca. 20 mL and, on cooling to room temperature, the desired compound precipitated. It was collected, washed with  $\text{CH}_2\text{Br}_2$  and then hexane and dried under vacuum to yield 3.33 g (63%) of colorless crystals, mp

236°C (decomp.). Anal. Calcd. for  $C_{14}H_{24}B_2Br_2N_4$  (429.80): C, 39.12; H, 5.63; B, 5.03; Br, 37.18; N, 13.04. Found: C, 39.21; H, 5.72; B, 4.69; Br, 37.31; N, 13.25.

NMR data (solution in  $CDCl_3$ ):  $\delta(^1H)$  = 6.20 (1H, s), 2.84 (3H, s), 2.48 (3H, s), 0.82 (2H, q,  $J = 7.7$ ), 0.34 (3H, t,  $J = 7.6$ );  $\delta(^{11}B)$  = +4.8 (1B,  $h_{\frac{1}{2}} = 200$  Hz), -6.7 (1B,  $h_{\frac{1}{2}} = 30$  Hz).

1,3,5,7-Tetramethyl-4,4,8,8-tetraethylpyrazabole.<sup>7</sup> NMR data (solution in  $CDCl_3$ ):  $\delta(^1H)$  = 6.00 (1H, s), 2.41 (6H, s), 0.77 (4H, q,  $J = 7.5$ ), 0.39 (6H, t,  $J = 7.5$ );  $\delta(^{11}B)$  = 3.5 ( $h_{\frac{1}{2}} = 150$  Hz);  $\delta(^{13}C)$  = 145.4 (s), 111.8 (d,  $J = 175$ ), 16.4\*, 15.4 (q,  $J = 129$ ), 10.2 (q,  $J = 124$ ). Literature:<sup>7</sup>  $\delta(^1H)$  = 5.70 (1H, s), 2.40 (6H, s), 0.92-0.25 (m);  $\delta(^{11}B)$  = 4.4.

1,3,5,7-Tetramethyl-4,8-dibromopyrazabole. A solution of 5.4 g (25 mmole) of 1,3,5,7-tetramethylpyrazabole<sup>7</sup> in 150 mL  $CH_2Br_2$  was added dropwise with stirring to a solution of 50 g (0.2 mole) of  $BBr_3$  in 50 mL  $CH_2Br_2$ . A precipitate formed immediately with slight warming of the reaction mixture. The mixture was stirred for 3 h at ambient temperature and the precipitate was collected. It was washed with  $CH_2Br_2$  and then a small amount of  $CHCl_3$  and dried under vacuum to give 7.9 g (84.5%) of the moisture-sensitive title compound, mp 298-305°C (with slight decomp). Anal. Calcd. for  $C_{10}H_{16}B_2Br_2N_4$  (373.70): C, 32.14; H, 4.32; B, 5.79; Br, 42.76; N, 14.99. Found: C, 31.70; H, 4.32; B, 6.15; Br, 43.00; N, 14.84.

NMR data (solution in  $CDCl_3$ ):  $\delta(^1H)$  = 6.11(s) + 5.84(s) (1H, ratio 2:1), 2.54(s) + 2.38(s) (6H, ratio 2:1);  $\delta(^{11}B)$

(proton-decoupled) =  $-8.3^*$ ,  $-10.3^*$  ( $h_{\frac{1}{2}}$  ca. 600 Hz);  $\delta(^{13}\text{C})$  (proton-decoupled) = (148.6), 145.0, 106.5, 93.2, (12.6), (12.1), 11.0. - IR:  $\nu(\text{BH}) = 2510 \text{ cm}^{-1}$ . - Mass spectrum (20 eV, probe temperature:  $180^\circ\text{C}$ ):  $M/z$  (relative abundance) = 296 (10), 295 (94), 294 (58), 293 (100), 292 (48), 291 (6), 214 (10), 213 (71), 212 (35), 211 (6). In addition, numerous peaks of lower than 5% relative abundances were observed. In particular, these include a low-intensity ion cluster in the parent ion region ( $M/z = 376$  to 369).

1,3,5,7-Tetramethyl-4,4,8,8-tetrabromopyrazabole. A solution of 12 mL  $\text{Br}_2$  in 50 mL  $\text{CH}_2\text{Br}_2$  was added dropwise with stirring to a solution of 10.8 g (50 mmol) of 1,3,5,7-tetramethylpyrazabole<sup>7</sup> in 300 mL  $\text{CH}_2\text{Br}_2$ . The mixture was refluxed for 3 h and was then reduced to 50 mL volume by distillation. After cooling to room temperature, the precipitate was collected, washed with  $\text{CH}_2\text{Br}_2$  and then with petroleum ether. The slightly yellow crystals were slurried with 15 mL  $\text{CHCl}_3$ , filtered and dried to give 11.0 g (83%) of colorless crystals, mp  $336\text{--}338^\circ\text{C}$  (decomp). Calcd. for  $\text{C}_{10}\text{H}_{14}\text{Br}_4\text{N}_4$  (531.49): C, 22.60; H, 2.65; Br, 4.07; N, 10.54. Found: C, 22.62; H, 2.47; Br, 3.92; N, 10.74.

NMR data (solution in  $\text{CDCl}_3$ ):  $\delta(^1\text{H}) = 6.31$  (s, 1H), 2.87 (s, 6H);  $\delta(^{11}\text{B}) = -7.5$  ( $h_{\frac{1}{2}} = 30$  Hz);  $\delta(^{13}\text{C}) = 152.1$  (s), 114.3 (d,  $J = 182$ ), 15.8 (q,  $J = 131$ ). - Mass spectrum:  $M/z$  (relative abundance) = 455 (31), 454 (25), 453 (95), 452 (57), 451 (100), 450 (49), 449 (37), 448 (16), 293 (21), 292 (27), 291 (32), 290 (27), 289 (11), 187 (8), 186.5 (5), 186 (15),

185.5 (15), 185 (9), 117 (5). Numerous additional ion peaks of lower than 5% relative abundances include a parent ion cluster  $M/z = 536$  to 526.

1,3,5,7-Tetramethyl-4,4,8,8-tetrakis(pyrazol-1'-yl)pyrazabole.

A mixture of 5.32 g (10 mmole) of 1,3,5,7-tetramethyl-4,4,8,8-tetrabromopyrazabole, 5.30 g (12.5 mmole) of potassium pyrazolyl-1-ate and 125 mL toluene was refluxed with stirring for 50 h. The hot mixture was filtered and the residue was washed with hot toluene. Solvent was evaporated from the combined toluene solutions. The crystalline residue was dissolved in chloroform and the solvent was evaporated from the clear filtrate. The solid residue was redissolved in 30 mL hot toluene. On standing (and slow evaporation of the solvent), several crystalline fractions were collected. The first one was essentially pure compound which was further purified by sublimation (material subliming under vacuum and at a bath temperature up to 150°C was discarded) to give ca. 2 g of product, mp 235-240°C. Based on mass spectral data the product contained a very minor impurity of  $B_2pz_3pz'_3$  ( $M/z$  508) and even  $B_2pz_2pz'_4$  ( $M/z$  536) which could, however, not be detected in the  $^1H$  NMR spectrum. Anal. Calcd. for  $C_{22}H_{26}B_2N_{12}$  (480.16): C, 55.03; H, 5.46; B, 4.50; N, 35.01. Found: C, 54.61; H, 5.63; B, 4.56; N, 35.06.

NMR data (solution in  $CDCl_3$ ):  $\delta(^1H)$  = 7.63 (2H, d,  $J = 1.45$ ), 6.65 (2H, d,  $J = 2.5$ ), 6.10 (1H, s), 6.04 (2H, unsymmetrical t = 2 overlapping d), 1.63 (6H, s);  $\delta(^{11}B)$  = -0.2 ( $h_z = 40$  Hz);  $\delta(^{13}C)$  (proton-decoupled) = 151.3, 142.6, 133.9, 112.2, 105.8, 11.3.

4,4,8,8-Tetrakis(pyrazol-1'-yl)pyrazabole.<sup>12</sup> NMR data (solution in  $\text{CD}_3\text{CN}$ ):  $\delta(^1\text{H}) = 7.57$  (2H, d,  $J = 2.3$ ),  $7.51$  (2H, d,  $J = 1.5$ ),  $6.75$  (2H, d,  $J = 2.2$ ),  $6.65$  (1H, t,  $J = 2.4$ ),  $6.07$  (2H, t,  $J = 1.7$ );  $\delta(^{11}\text{B}) = 0.7$  ( $h_{\frac{1}{2}} = 35$  Hz);  $\delta(^{13}\text{C})$  (proton-decoupled) =  $143.6$ ,  $141.1$ ,  $134.9$ ,  $109.5$ ,  $107.0$ . In  $\text{DMSO}-d_6$ :  $\delta(^{11}\text{B}) = 0.2$  ( $h_{\frac{1}{2}} = 120$  Hz). Additional NMR data have been reported elsewhere.<sup>6</sup> - Infrared spectrum (KBr pellet):  $3100$  (sh),  $3090$  (w),  $1770$  (vw),  $1721$  (vw),  $1507$  (m),  $1493$  (sh),  $1411$  (s, b),  $1382$  (vs),  $1342$  (m),  $1326$  (ms),  $1291$  (vs, b),  $1247$  (sh),  $1233$  (s),  $1219$  (s),  $1213$  (s),  $1202$  (s),  $1186$  (m),  $1143$  (m),  $1095$  (s),  $1082$  (vs),  $1060$  (m),  $1045$  (m),  $1037$  (m),  $1023$  (m),  $971$  (vw),  $946$  (sh),  $943$  (m),  $933$  (vw),  $918$  (m),  $889$  (sh),  $883$  (sh),  $873$  (s),  $859$  (sh),  $851$  (sh),  $838$  (sh),  $833$  (vs),  $816$  (m),  $811$  (s),  $786$  (s),  $782$  (sh),  $772$  (s),  $760$  (sh),  $755$  (vs),  $673$  (sh),  $668$  (w),  $658$  (w),  $650$  (vw).

1,3,5,7-Tetramethyl-4,8-bis(3',5'-dimethylpyrazol-1'-yl)-pyrazabole, second modification. A mixture of  $20.2$  g ( $0.05$  mole) of 1,3,5,7-tetramethylpyrazabole and  $9.6$  g ( $0.1$  mole) of 3,5-dimethylpyrazole was heated in an oil-bath of  $180^\circ\text{C}$  for 12 h and the temperature was then slowly increased to  $220^\circ\text{C}$  for 2 h. The melt now contained a considerable amount of precipitate and was cooled to room temperature. The product was washed with copious amounts of hot benzene and approximately 6 g of a powdery material remained. It did not melt at temperatures up to  $350^\circ\text{C}$ , where it began to decompose. The elemental analysis as well as NMR and mass spectral data of the material were identical to



those of the previously described 1,3,5,7-tetramethyl-4,8-bis-(3',5'-dimethylpyrazol-1'-yl)pyrazabole,<sup>5</sup> although this latter material had a sharp mp 286-288°C. However, the infrared spectra of the two modifications showed several significant differences (see below).


In an alternate experiment, one molar equivalent of the previously described<sup>5</sup> 1,3,5,7-tetramethyl-4,8-bis(3',5'-dimethylpyrazol-1'-yl)pyrazabole was heated for 8 h with two molar equivalents of 3,5-dimethylpyrazole to 180°C. At that time, again a considerable amount of precipitate had formed and the reaction mixture was worked up as above. The benzene-insoluble portion showed an identical IR spectrum to the one of the second modification of the title compound as prepared above, whereas the <sup>1</sup>H NMR spectrum was again identical to that of the previously described<sup>5</sup> modification. After evaporation of the solvent from the NMR solution, the remaining crystals had the mp 286-288°C, i.e., that of the normal modification.

IR spectral data: Previously described<sup>5</sup> modification (KBr pellet, 3200 to 600 cm<sup>-1</sup> region only): 3218(m), 3082(w), 2980(sh), 2960(m), 2922(ms), 2900(sh), 2860(w), 2735(w), 2631(vw), 2462(vs), 2355(w), 2220(w), 2130(vw), 1625(w, br), 1562(sh), 1550 (vs), 1539(m), 1502(w), 1460(sh), 1452/1435/1415 (vs), 1390(vs), 1374(sh), 1368(w), 1348/1338(vs), 1264(s), 1204(sh), 1198(vs), 1155(vs), 1130(sh), 1070(vs), 1050(m), 1038(wm), 1020(wm), 1008(ms), 975(sh), 905(sh), 898(s), 812(vs), 802(s), 765(sh), 758(m), 685(w), 660 (m), 640(s). The spectrum of the second modification as prepared above was essentially identical to the preceding data, with the exception of the

following differences: The  $\nu(\text{BH})$  band had now shifted from  $2462\text{ cm}^{-1}$  to  $2495\text{ cm}^{-1}$ ; the medium-intensity band at  $975\text{ cm}^{-1}$  had become a very weak peak but a new band appeared at  $852\text{ cm}^{-1}$  (m); the very strong absorption at  $812\text{ cm}^{-1}$  had shifted to  $770\text{ cm}^{-1}$ ; and the  $660\text{ cm}^{-1}$  band (m) was missing from the spectrum of the second modification.

## Results and Discussion


Preparation and Characterization of Unsymmetrically Substituted Pyrazaboles. Recently,<sup>5</sup> the reaction of  $\text{K}[\text{Bpz}_4]$  (pz =  $\text{N}_2\text{C}_3\text{H}_3$  = pyrazolyl) with  $(\text{CH}_3)_3\text{NBH}_2\text{I}$  has been described as a new approach for the preparation of unsymmetrically substituted pyrazaboles of the type  $\text{R}_2\text{B}(\mu\text{-pz})_2\text{BR}'_2$ . The basic reaction is not limited to the cited case as is now documented by the reaction of  $\text{K}[\text{R}_2\text{Bpz}_2]$  ( $\text{R} = \text{C}_6\text{H}_5$ ) with  $(\text{CH}_3)_3\text{NBH}_2\text{I}$  according to eq (1).



The  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra of the resultant 4,4-diphenylpyrazabole were readily assigned on the basis of the observed fine structure, coupling constants, HETCOR and HOMCOR-2D NMR experiments, and relevant NMR data of the two related compounds  $\text{H}_2\text{B}(\mu\text{-pz})_2\text{BH}_2$  and  $(\text{C}_6\text{H}_5)_2\text{B}(\mu\text{-pz})_2\text{B}(\text{C}_6\text{H}_5)_2$ . Thus the  $\delta(^1\text{H})/\delta(^{13}\text{C})$  signals 7.23/127.3, 6.97/133.6, and 7.23/126.8 ppm are those of the o-, m- and p-CH moieties, respectively, of the phenyl groups;

whereas those at 7.66/135.7 ppm are those of the pyrazoyl-CH closest to the  $B(C_6H_5)_2$  group ( $\delta(^{11}B) = +1.8$  ppm) and those at 7.37/136.3 ppm of the ones closest to the  $BH_2$  group ( $\delta(^{11}B) = -8.6$  ppm). The  $\delta(^1H)/\delta(^{13}C)$  signals of the central CH unit of the pyrazolyl rings are at 6.31/105.3 ppm.

4,4-Diphenylpyrazabole reacts with elemental bromine to yield 4,4-diphenyl-8,8-dibromopyrazabole according to eq (2).



The reaction proceeds in good yield and neither of the aromatic rings is attacked by  $Br_2$  or  $HBr$ .

In a corresponding reaction, 1,3,5,7-tetramethyl-4,4-diethylpyrazabole was converted to 1,3,5,7-tetramethyl-4,4-diethyl-8,8-dibromopyrazabole in good yield. Thus, in contrast to the reaction with  $BBr_3$ ,<sup>4</sup> reaction with  $Br_2$  displaces only boron-bonded hydrogen but no organic substituents. However, it should be noted that, based on mass spectral and  $^1H$  NMR data of the crude reaction products, small amounts of  $R_2B(\mu-pz')_2BR_2$  ( $R = C_2H_5$ ) and  $Br_2B(\mu-pz')_2BBr_2$  were also formed. This observation suggests at least some opening of the central  $B_2N_4$  ring of the original pyrazabole during the course of the reaction!

As based on mass spectral and  $^1H$  NMR data, the reaction of  $K[R_2Bpz_2]$  ( $R = C_6H_5$ ) with two molar equivalents of  $(C_2H_5)_2OBF_3$  gave some of the expected  $R_2B(\mu-pz)_2BF_2$ . However, a small amount of  $F_2B(\mu-pz)_2BF_2$  was also observed and  $R_2B(\mu-pz)_2BR_2$  was the major product; only the latter

could be isolated in pure state! Also, the reaction of  $K[Bpz_4]$  with  $R_2BBr$  did not yield the desired  $R_2B(\mu-pz)_2Bpz_2$ ; rather, as based on  $^1H$  NMR data, a mixture was obtained containing at least two different pyrazaboles. Furthermore, all attempts to prepare  $pz_2B(\mu-pz)_2BF_2$  from the interaction of  $K[Bpz_4]$  with  $(C_2H_5)_2OBF_3$  failed. The only identified product of this latter reaction was  $F_2B(\mu-pz)_2BF_2$ .

These cited observations suggest a fairly complex path for the interaction of poly(pyrazol-1-yl)borates with trigonal boranes or Lewis base adducts thereof. This is in consonance with the previous<sup>5</sup> report that even in the reaction of  $K[Bpz_4]$  with  $(CH_3)_3NBH_2I$  substantial quantities of  $H_2B(\mu-pz)_2BH_2$  were formed. Hence, the few observed cases to yield the desired products in reasonable quantity<sup>5,11</sup> seem to be fortuitous and a more detailed study is mandated before final conclusions about the general utility of the interaction of poly(pyrazol-1-yl)-borates with trigonal boranes for the formation of pyrazaboles can be drawn.

Chemical Studies on 1,3,5,7-Tetramethylpyrazaboles. The condensation of 3,5-dimethylpyrazole,  $Hpz'$ , with trimethylamine-borane proceeds readily to yield  $H_2B(\mu-pz')_2BH_2$ .<sup>7</sup> The latter compound reacts with additional  $Hpz'$  to yield  $Hpz'B(\mu-pz')_2BHpz'$  in a smooth reaction, but subsequent formation of  $pz'_2B(\mu-pz')_2Bpz'_2$  requires excessive reaction times at high temperatures.<sup>5</sup> Even when a clear melt of  $Hpz'B(\mu-pz')_2BHpz'$  and  $Hpz'$  was kept at temperatures near  $180^\circ C$  for six hours,


no formation of  $\text{pz}'_2\text{B}(\mu\text{-pz}')_2\text{Bpz}'_2$  was observed as indicated by a mass spectrum of the product. However, at that time some solid material slowly precipitated from the melt and the amount of precipitate increased within two more hours of heating. In contrast to the other components of the reaction mixture, this precipitate was essentially insoluble in hot benzene and could thus be separated. It was identified by elemental analysis as well as NMR and mass spectroscopic data as  $\text{Hpz}'\text{B}(\mu\text{-pz}')_2\text{BHpz}'$ . However, although the cited analytical data were in complete agreement with those of the previously described material,<sup>5</sup> the melting behavior was distinctly different: It did not melt at temperatures up to  $350^\circ\text{C}$  where decomposition began. This contrasts with the normal modification of the compound which melts at  $286\text{-}288^\circ\text{C}$  without decomposition. Also, there were few but significant differences in the infrared spectra of the two materials. It should be noted that on evaporation of a solution of the high melting material (which was used for the NMR spectroscopic studies), the resultant crystalline residue showed the melting point of the previously known modification of the compound, i.e.,  $286\text{-}288^\circ\text{C}$ .

Based on these observations, the high melting material is assumed to be a second modification of  $\text{Hpz}'\text{B}(\mu\text{-pz}')_2\text{BHpz}'$ . It is noteworthy that only when this second modification has been formed, reaction with additional  $\text{Hpz}'$  initiates and slowly leads to the formation of  $\text{pz}'_2\text{B}(\mu\text{-pz}')_2\text{Bpz}'_2$  (as indicated by mass spectral data). This observation suggests that the

boron-bonded hydrogen atoms of the normal modification of  $\text{Hpz}'\text{B}(\mu\text{-pz}')_2\text{BHpz}'$  are much less reactive than those of the second modification.

The crystal structure of  $\text{Hpz}'\text{B}(\mu\text{-pz}')_2\text{BHpz}'$  has been shown to involve a  $\text{B}_2\text{N}_4$  ring in chair conformation and the two terminal pyrazolyl groups were found to be in trans configuration and in axial position.<sup>13</sup> (Presumably, the sample involved was that of the previously described<sup>5</sup> modification of mp 286-288°C; unfortunately, no crystals suitable for a X-ray structure determination could be obtained from the second modification as described above.) Thus, the remaining B-H bonds of the molecule are in equatorial position. In this context it is of interest to note that the axial B-H bonds of  $\text{H}_2\text{B}(\mu\text{-pz})_2\text{BH}_2$ <sup>14</sup> and  $\text{H}_2\text{B}(\mu\text{-pz})_2\text{Bpz}_2$ <sup>15</sup> have been found to be longer, and hence weaker, than the equatorial ones. This difference in bond length = reactivity seems to be also borne out by the interaction of  $\text{H}_2\text{B}(\mu\text{-pz}')_2\text{BH}_2$  with  $\text{BBr}_3$  (see below).

Considering all of the preceding data it appears that the second modification of  $\text{Hpz}'\text{B}(\mu\text{-pz}')_2\text{BHpz}'$  is macromolecular in nature. This could be explained by assuming that one of the bridging B-N bonds of the pyrazabole is cleaved on prolonged heating to high temperatures to yield the structural unit 2.



2

The species then could exist in a cyclic arrangement of several such units or even in a linear polymer containing trigonal boron in chain-terminating groups. In any case, the macromolecular nature would explain the lack of melting prior to thermal decomposition. It would also explain the differences in the infrared spectra of the two modifications and, of course, would influence the reactivity of the B-H bonds: The latter appear to be more reactive (in a condensation process) in the polymer. The identity of the NMR spectra of both modifications suggests that, once in solution, the macromolecule breaks down to the normal pyrazaborole structure. (Note: Indeed, the polymer is considerably less readily soluble in  $\text{CHCl}_3$  as compared to the normal modification of the compound.)

In a previous study it has been shown that the condensation of pyrazaborole with pyrazoles at high temperatures must involve

at least partial cleavage of the central  $B_2N_4$  ring of the pyrazabole.<sup>5</sup> The present findings seem to be in consonance with that observation. Apparently, 4,8-disubstitution at  $BH_2$  sites of pyrazaboles occurs quite readily in a condensation process; however, a second such step may more readily proceed through polymeric intermediates formed by ring opening, although these have not been observed previously. This interpretation would readily account for the product mixture which was obtained on reacting  $H_2B(\mu\text{-pz})_2BH_2$  with C-substituted pyrazoles.<sup>5</sup>

In this connection it is of interest to note that the boat conformation of the central  $B_2N_4$  ring seems to be the preferred structural arrangement of pyrazaboles.<sup>4</sup> It seems possible that, even in low-temperature electrophilic-induced substitution processes, the arrangement of terminal substituents at the boron sites of pyrazaboles governs the reactivity to a major extent. This is already suggested by the pattern of B-disubstitution of a preformed pyrazabole, which always leads to a 4,8-disubstituted product rather than a geminal species. Furthermore, it is apparent that C-substitution at bridging pyrazolyl groups of pyrazaboles also induces specific features. For example, the parent pyrazabole reacts readily with an excess of either  $BBr_3$  or  $Br_2$  to form  $Br_2B(\mu\text{-pz})_2BBr_2$ .<sup>8</sup> In contrast, 1,3,5,7-tetramethylpyrazabole reacts with excess  $BBr_3$  to form only the corresponding 4,8-dibromo derivative. The latter exists as a mixture of cis and trans isomer, as was



the case with the corresponding derivative of the parent compound, i.e.,  $\text{HBrB}(\mu\text{-pz})_2\text{BBr}$ .<sup>14</sup> On the other hand, reaction of 1,3,5,7-tetramethylpyrazabole with excess  $\text{Br}_2$  affords the 4,4,8,8-tetrabromo derivative,  $\text{Br}_2\text{B}(\mu\text{-pz}')_2\text{BBr}_2$ , in essentially quantitative yield.

In another experiment, Kpz was reacted with  $\text{Br}_2\text{B}(\mu\text{-pz}')_2\text{BBr}_2$  in refluxing toluene to yield  $\text{pz}_2\text{B}(\mu\text{-pz}')_2\text{Bpz}_2$ . This latter compound is the first example of an isolated and characterized pyrazolylpyrazabole where a boron atom is bonded to two chemically different pyrazole moieties. Remarkably, mass spectral data on the crude reaction product indicated the formation of the pyrazaboles of the composition  $\text{B}_2\text{pz}_3\text{pz}'_3$  and  $\text{B}_2\text{pz}_2\text{pz}'_4$  as minor byproducts. Their formation suggests again opening of the central  $\text{B}_2\text{N}_4$  ring of the starting material during the course of the reaction. Even though this seems to occur only to a small extent, it is apparent that such pyrazabole-ring opening may be much more common than heretofore assumed. This is in consonance with the various observations described above.

The observed  $\delta(^1\text{H})/\delta(^{13}\text{C})$  data of  $\text{pz}_2\text{B}(\mu\text{-pz}')_2\text{Bpz}_2$  are readily assigned on comparison with those of  $\text{pz}_2\text{B}(\mu\text{-pz})_2\text{Bpz}_2$ .<sup>6</sup> The shifts 7.63/142.6, 6.04/105.8, and 6.65/133.9 of the former belong to the 3, 4, and 5 position, respectively, of the terminal pyrazolyl groups. They are essentially identical to those of the terminal pyrazolyl groups in  $\text{pz}_2\text{B}(\mu\text{-pz})_2\text{Bpz}_2$  with 7.70/142.6, 6.16/105.7, and 6.81/133.6, respectively. Also, the values for the 2,6 positions of the pyrazabole

skeleton of  $\text{pz}_2\text{B}(\mu\text{-pz}')_2\text{Bpz}_2$  with 6.10/112.2 compare well with the corresponding data<sup>5</sup> of  $\text{pz}_2'\text{B}(\mu\text{-pz}')_2\text{Bpz}_2'\cdot\text{H}_2\text{O}$  with 5.91/111.7 (and 5.88/111.1 in the anhydrous compound).

The compound  $\text{pz}_2'\text{B}(\mu\text{-pz}')_2\text{Bpz}_2'\cdot\text{H}_2\text{O}$  can be dehydrated, though with some difficulty.<sup>5</sup> The anhydrous material is readily reconverted to the monohydrate by stirring the compound with water for a few minutes. However, anhydrous  $\text{pz}_2'\text{B}(\mu\text{-pz}')_2\text{Bpz}_2'$  does not interact with liquid anhydrous ammonia, not even at room temperature in a sealed tube. Also, no reaction occurred when a solution of the anhydrous compound in  $\text{CHCl}_3$  was mixed with a saturated solution of  $\text{NH}_3$  in the same solvent and the mixture was subsequently evaporated. The starting material was recovered unchanged in all cases.

Acknowledgment. This work was supported by the Office of Naval Research (K.N.). One of us (P.M.N.) gratefully acknowledges an Ashland Oil Summer Research Fellowship.

#### References

- (1) Boron-Nitrogen Compounds. 101 (K.N.). For part 100 of this series see ref. 14.
- (2) "Gmelin Handbuch der Anorganischen Chemie"; Springer-Verlag: West Berlin 1975; Vol. 23, Supplement Boron Compounds 5, pp. 1-8.
- (3) Weber, W.; Niedenzu, K. J. Organomet. Chem., 1981, 205, 147.
- (4) Niedenzu, K.; Noeth, H. Chem. Ber., 1983, 116, 1132.
- (5) Niedenzu, K.; Niedenzu, P. M. Inorg. Chem., 1984, 23, 0000.

- (6) Layton, W. J.; Niedenzu, K.; Smith, S. L. Z. Anorg. Allg. Chem., 1982, 495, 52.
- (7) Trofimenko, S. J. Am. Chem. Soc., 1967, 89, 3165.
- (8) Ryschkewitsch, G. E.; Wiggins, J. W. Inorg. Synth., 1972, 12, 116.
- (9) Myers, W. H.; Ryschkewitsch, G. E.; Matnur, W. A.; King, R. W. Inorg. Chem., 1975, 14, 2874.
- (10) Denniston, M. L.; Chiusano, M.; Brown, J.; Martin, D. R. J. Inorg. Nucl. Chem., 1976, 38, 379.
- (11) Trofimenko, S. Inorg. Chem., 1969, 8, 1714.
- (12) Trofimenko, S. J. Am. Chem. Soc., 1967, 89, 4948.
- (13) Alcock, N. W.; Sawyer, J. F. Acta Crystallogr., 1974, B30, 2899.
- (14) Hanecker, E.; Hodgkins, T. G.; Niedenzu, K.; Noeth, H. Inorg. Chem., in hand of editor.
- (15) Brock, C. P.; Hanecker, E.; Niedenzu, K.; Noeth, H. to be published.

**END**

**FILMED**

**1-85**

**DTIC**